

University of Wisconsin - Madison

MADPH-98-1050

April 1998

The AMANDA Neutrino Telescope and the Indirect Search for Dark Matter*

Francis Halzen, for the AMANDA Collaboration

R.C. Bay, Y. He, D. Lowder, P. Miocinovic, P.B. Price, M. Solarz, K. Woschnagg

University of California, Berkeley, USA

S.W. Barwick, Je. Booth, P.C. Mock, R. Porrata, E. Schneider, G. Yodh

University of California, Irvine, USA

D. Cowen

University of Pennsylvania, USA

M. Carlson, C.G.S. Costa, T. DeYoung L. Gray, F. Halzen, R. Hardtke, J. Jacobsen,

V. Kankhadai, A. Karle, I. Liubarsky, R. Morse, S. Tilav

University of Wisconsin, Madison, USA

T.C. Miller

Bartol Research Institute, USA

E.C. Andrés, P. Askebjør, L. Bergström, A. Bouchta, E. Dalberg, P. Ekström,

A. Goobar, P.O. Hulth, C. Walck

Stockholm University, Sweden

A. Hallgren, C. P. de los Heros, P. Marciniwski, H. Rubinstein

University of Uppsala, Sweden

S. Carius, P. Lindahl

Kalmar University, Sweden

A. Biron, S. Hundertmark, M. Leuthold, P. Niessen, C. Spiering, O. Streicher,

T. Thon, C.H. Wiebusch, R. Wischnewski

DESY — Institute for High Energy Physics, Germany

D. Nygren

Lawrence Berkeley National Laboratory, USA

A. Jones, S. Hart, D. Potter, G. Hill, R. Schwarz

South Pole Winter-Overs, Antarctica

Abstract

With an effective telescope area of order 10^4 m^2 , a threshold of $\sim 50 \text{ GeV}$ and a pointing accuracy of 2.5 degrees, the AMANDA detector represents the first of a new generation of high energy neutrino telescopes, reaching a scale envisaged over 25 years ago. We describe its performance, focussing on the capability to detect halo dark matter particles via their annihilation into neutrinos.

*Talk presented at the *3rd International Symposium on Sources and Detection of Dark Matter in the Universe (DM98)*, Santa Monica, California, Feb. 1998.

1 The Indirect Detection of Halo Dark Matter

High energy neutrino telescopes are multi-purpose instruments; their science mission covers particle physics, astronomy and astrophysics, cosmology and cosmic ray physics. Their deployment creates new opportunities for glaciology and oceanography, possibly geology of the earth's core[1]. Astronomy with neutrinos does have definite advantages. They can reach us, essentially without attenuation in flux, from the largest red-shifts. The sky is, in contrast, partially opaque to high energy photons and protons because of energy-loss suffered in interactions with infrared light, CMBR photons and radio waves[2]. They do not reach us from distances much larger than tens of megaparsecs once their energy exceeds thresholds of 10 TeV for photons and 5×10^7 TeV for protons. (Below this energy charged protons do not point back to their sources.) The drawback is that neutrinos are difficult to detect: the small interaction cross sections that enable them to travel without attenuation over a Hubble radius, are also the reason why kilometer-scale detectors are required in order to capture them in sufficient numbers to do astronomy[3]. Some opportunities may, however, be unique. If, for instance, the sources of the highest energy cosmic rays are beyond 10^2 Mpc, conventional astronomy is unlikely to discover them.

Some science missions do not require a detector of kilometer size. The best opportunities to search for halo dark matter are, in fact, associated with the present instrument which, while smaller in telescope area than the planned extension of AMANDA to ICE3 (ICECUBE), has a lower threshold.

At this meeting, the capability of neutrino telescopes to discover the particles that constitute the dominant, cold component of the dark matter is of special interest. The existence of the weakly interacting massive particles (WIMPs) is inferred from observation of their annihilation products. Cold dark matter particles annihilate into neutrinos; *massive* ones will annihilate into *high-energy* neutrinos which can be detected in high-energy neutrino telescopes. This so-called indirect detection is greatly facilitated by the fact that the earth and the sun represent dense, nearby sources of accumulated cold dark matter particles. Galactic WIMPs, scattering off nuclei in the sun, lose energy. They may fall below escape velocity and be gravitationally trapped. Trapped WIMPs eventually come to equilibrium and accumulate near the center of the sun. While the WIMP density builds up, their annihilation rate into lighter particles increases until equilibrium is achieved where the annihilation rate equals half of the capture rate. The sun has thus become a reservoir of WIMPs which we expect to annihilate mostly into heavy quarks and, for the heavier WIMPs, into weak bosons. The leptonic decays of the heavy quark and weak boson annihilation products turn the sun and earth into nearby sources of high-energy neutrinos with energies in the GeV to TeV range. Figure 1 displays the neutrino flux from the center of the earth calculated in the context of supersymmetric dark matter theories[4]. The direct capture rate of the WIMPs in germanium detectors is shown for comparison. Contours indicate the parameter space favored by grand unified theories. Most of this parameter space can be covered by improving the capabilities of existing detectors by 2 orders of magnitude. Existing neutrino detectors have already excluded fluxes of neutrinos from the earth's center of order 1 event per 1000 m² per year. The best limits have been obtained by the Baksan experiment[5]. They are already excluding relevant parameter space of supersymmetric models. We will show that, with data already on tape, the AMANDA detector will have an unmatched discovery reach for WIMP masses in excess of 100 GeV.




Figure 1: Direct and indirect detection rates (for neutrinos from the center of the earth in the figure shown) of cold dark matter particles predicted by supersymmetric theory. Grand unified theories favor the parameter space indicated. Part of it is already excluded by present experiments as indicated by the horizontal and vertical lines.

The potential of neutrino telescopes as dark matter detectors has been documented in detail[6]. With a sensitivity which increases with the WIMP mass, they are complementary to direct, cryogenic detectors. They can detect WIMPS beyond the kinematic limits of the LHC: about 500 GeV for neutralinos. A striking way to illustrate their potential is to use the possible detection[7] in the DAMA NaI detector in the Gran Sasso tunnel as an example. If their seasonal variation is indeed evidence for WIMPS, observation of a signal in an exposure of 4500 kg days requires a WIMP-nucleon cross section of $10^{-42} \sim 10^{-41}$ cm² for a WIMP mass of 50~150 GeV. This information is sufficient to calculate their trapping and annihilation rate in the sun and earth. Both will be a source of, on average, 100 neutrinos per year of WIMP origin in the existing AMANDA detector with an effective area of 10⁴ m². The exact rate varies with the mass of the WIMP.

2 Status of the AMANDA Project

First generation neutrino detectors, launched by the bold decision of the DUMAND collaboration over 20 years ago to construct such an instrument, are designed to reach a relatively large telescope area and detection volume for a neutrino threshold of tens of GeV, not higher. This relatively low threshold permits calibration of the novel instrument on the known flux of atmospheric neutrinos. Its architecture is optimized for reconstructing the Cherenkov light front radiated by an up-going, neutrino-induced muon. Up-going muons are to be identified in a background of down-going, cosmic ray muons which are more than 10⁵ times more frequent for a depth of 1~2 kilometers. The method is sketched in Fig. 2.




Figure 2: The arrival times of the Cherenkov photons in 6 optical modules determine the direction of the muon track.

Construction of the first-generation AMANDA detector[8] was completed in the austral summer 96–97. It consists of 300 optical modules deployed at a depth of 1500–2000 m; see Fig. 3. An optical module (OM) consists of an 8 inch photomultiplier tube and nothing else. OM’s have only failed when the ice refreezes, at a rate of less than 3 percent. Calibration of this detector is in progress, although data has been taken with 80 OM’s which were deployed one year earlier in order to verify the optical properties of the ice (AMANDA-80).

As anticipated from transparency measurements performed with the shallow strings[11] (see Fig. 3), we found that ice is bubble-free at 1400–1500 meters and below. The performance of the AMANDA detector is encapsulated in the event shown in Fig. 4. Coincident events between AMANDA-80 and four shallow strings with 80 OM’s have been triggered for one year at a rate of 0.1 Hz. Every 10 seconds a cosmic ray muon is tracked over 1.2 kilometers. The contrast in detector response between the strings near 1 and 2 km depths is dramatic: while the Cherenkov photons diffuse on remnant bubbles in the shallow ice, a straight track with velocity c is registered in the deeper ice. The optical quality of the deep ice can be assessed by viewing the OM signals from a single muon triggering 2 strings separated by 79.5 m; see Fig. 4b. The separation of the photons along the Cherenkov cone is well over 100 m, yet, despite some evidence of scattering, the speed-of-light propagation of the track can be readily identified.

The optical properties of the ice are quantified by studying the propagation in the ice of pulses of laser light of nanosecond duration. The arrival times of the photons after 20 m and 40 m are shown in Fig. 5 for the shallow and deep ice[9]. The distributions have been normalized to equal areas; in reality, the probability that a photon travels 70 m in the deep ice is $\sim 10^7$ times larger. There is no diffusion resulting in loss of information on the geometry of the Cherenkov cone in the deep bubble-free ice.




Figure 3: The Antarctic Muon And Neutrino Detector Array (AMANDA).

3 AMANDA: before and after

The AMANDA detector was antecedently proposed on the premise that inferior properties of ice as a particle detector with respect to water could be compensated by additional optical modules. The technique was supposed to be a factor 5~10 more cost-effective and, therefore, competitive. The design was based on then current information[10]:

- the absorption length at 370 nm, the wavelength where photomultipliers are maximally efficient, had been measured to be 8 m;




Figure 4a: Cosmic ray muon track triggered by both shallow and deep AMANDA OM's. Trigger times of the optical modules are shown as a function of depth. The diagram shows the diffusion of the track by bubbles above 1 km depth. Early and late hits, not associated with the track, are photomultiplier noise.

- the scattering length was unknown;
- the AMANDA strategy would have been to use a large number of closely spaced OM's to overcome the short absorption length. Muon tracks triggering 6 or more OM's are reconstructed with degree accuracy. Taking data with a simple majority trigger of 6 OM's or more at 100 Hz yields an average effective area of 10^4 m^2 , somewhat smaller for atmospheric neutrinos and significantly larger for the high energy signals previously discussed.

The reality is that:

- the absorption length is 100 m or more, depending on depth[11];




Figure 4b: Cosmic ray muon track triggered by both shallow and deep AMANDA OM's. Trigger times are shown separately for each string in the deep detector. In this event the muon mostly triggers OM's on strings 1 and 4 which are separated by 79.5 m.

- the scattering length is 25~30 m (preliminary, this number represents an average value which may include the combined effects of deep ice and the refrozen ice disturbed by the hot water drilling);
- because of the large absorption length, OM spacings are similar, actually larger, than those of proposed water detectors. Also, a typical event triggers 20 OM's, not 6. Of these more than 5 photons are, on average, “not scattered.” A precise definition of “direct” photons will be given further on. In the end, reconstruction is therefore as before, although additional information can be extracted from scattered photons by minimizing a likelihood function which matches measured and expected delays[12].




Figure 5: Propagation of 510 nm photons indicate bubble-free ice below 1500 m, in contrast to ice with some remnant bubbles above 1.4 km.

The measured arrival directions of background cosmic ray muon tracks, reconstructed with 5 or more unscattered photons, are confronted with their known angular distribution in Fig. 6. There is an additional cut in Fig. 6 which simply requires that the track, reconstructed from timing information, actually traces the spatial positions of the OM's in the trigger. The power of this cut, especially for events distributed over only 4 strings, is very revealing. It can be shown that, in a kilometer-scale detector, geometrical track reconstruction using only the positions of triggered OM's is sufficient to achieve degree accuracy in zenith angle. We conclude from Fig. 6 that the agreement between data and Monte Carlo simulation is adequate. Less than one in 10^5 tracks is misreconstructed as originating below the detector[9]. Visual inspection reveals that the remaining misreconstructed tracks are mostly showers, radiated by muons or initiated by electron neutrinos, misreconstructed as up-going tracks of muon neutrino origin. At the 10^{-6} level of the background, candidate events can be identified; see Fig. 7. This exercise establishes that AMANDA-80 can be operated as a neutrino detector; misreconstructed showers can be readily eliminated on the basis of the additional information on the amplitude of OM signals. Monte Carlo simulation, based on this exercise, confirms that AMANDA-300 is a 10^4 m² detector with 2.5 degrees mean angular resolution[12]. We have verified the angular resolution of AMANDA-80 by reconstructing muon tracks registered in coincidence with a surface air shower array SPASE[13]. Figure 8 demonstrates that the zenith angle distribution of the coincident SPASE-AMANDA cosmic ray beam reconstructed by the surface array is quantitatively reproduced by reconstruction of the muons in AMANDA.




Figure 6: Reconstructed zenith angle distribution of muons triggering AMANDA-80: data and Monte Carlo. The relative normalization has not been adjusted at any level. The plot demonstrates a rejection of cosmic ray muons at a level of 10^{-5} .




Figure 7: A candidate up-going, neutrino-induced muon in the AMANDA-80 data. The numbers show the time sequence of triggered OMs, the size of the dots the relative amplitude of the signal.

4 Neutrinos from the earth's Center: AMANDA-80

We have performed a search[14] for upcoming neutrinos from the center of the earth. One should keep in mind that the preliminary results are obtained with only 80 OMs, incomplete calibration and only 6 months of data. We nevertheless obtain limits near the competitive level of less than 1 event per 250 m^2 per year for WIMP masses in excess of 100 GeV. With calibration of AMANDA-300 completed and 1 year of data on tape, we anticipate a sensitivity beyond the limit shown in Fig. 1. Increased sensitivity results from: lower threshold, better calibration (factor of 3), improved angular resolution (factor of ~ 2), longer exposure and, finally, an effective area larger by over one order of magnitude. Recall that, because the




Figure 8: Zenith angle distributions of cosmic rays triggering AMANDA and the surface air shower array SPASE. Reconstruction by AMANDA of underground muons agrees with the reconstruction of the air shower direction using the scintillator array.

search is limited by atmospheric neutrino background, sensitivity only grows as the square root of the effective area.

We reconstructed 6 months of filtered AMANDA-80 events subject to the conditions that 8 OM's report a signal in a time window of 2 microseconds. While the detector accumulated data at a rate of about 20 Hz, filtered events passed cuts [15] which indicate that time flows upwards through the detector. In collider experiments this would be referred to as a level 3 trigger. The narrow, long AMANDA-80 detector (which constitutes the 4 inner strings of AMANDA-300) thus achieves optimal efficiency for muons pointing back towards the center of the earth which travel vertically upwards through the detector. Because of edge effects the efficiency, which is, of course, a very strong function of detector size, is only a few percent after final cuts, even in the vertical direction. Nevertheless, we will identify background atmospheric neutrinos and establish meaningful limits on WIMP fluxes from the center of the earth.

That this data set, including prefiltering, is relatively well simulated by the Monte Carlo is shown in Fig. 9. The results reinforce the conclusions, first drawn from Figs. 6 and 8, that we understand the performance of the detector. Cuts are on the number of “direct” photons, i.e. photons which arrive within time residuals of $[-15; 25]$ ns relative to the predicted time. The latter is the time it takes for Cherenkov photons to reach the OM from




Figure 9: $\cos(\theta_{\text{rec}})$ is shown with cuts on the number of residuals in the interval $[-15; 25]$ ns. The histogram represents Monte Carlo simulations with a trigger of 8 or more hits in 2 msec and the dots represent the real data. (Notice that up and down directions are reversed from Fig. 6)

the reconstructed muon track. The choice of residual reflects the present resolution of our time measurements and allows for delays of slightly scattered photons. The reconstruction capability of AMANDA-80 is illustrated in Fig. 10. Comparison of the reconstructed zenith angle distribution of atmospheric muons and the Monte Carlo is shown in Fig. 10a for 3 cuts in N_{direct} . For $N_{\text{direct}} \geq 5$, the resolution is 2.2 degrees as shown in Fig. 10b.

The final cut selecting WIMP candidates requires 6 or more residuals in the interval $[-15, +15]$ ns and $\alpha \geq 0.1$ m/ns. Here α is the slope parameter obtained from a plane wave fit $z_i = \alpha t_i + \beta$, where z_i are the vertical coordinates of hit OMs and t_i the times at which they were hit. The two events surviving these cuts are shown in Fig. 11. Their properties are summarized in Table 1. The expected number of atmospheric neutrino events passing the same cuts is $4.8 \pm 0.8 \pm 1.1$. With only preliminary calibration, the systematic error in the time-calibration of the PMTs is ~ 15 ns. This reduces the number of expected events to $2.9 \pm 0.6 \pm 0.6$. The fact that the parameters of both events are not close to the cuts imposed, reinforces their significance as genuine neutrino candidates. Their large tracklengths suggest neutrino energies in the vicinity of 100 GeV which implies that the parent neutrino directions should align with the measured muon track to better than 2 degrees. Conservatively, we conclude that we observe 2 events on a background of 4.8 atmospheric neutrinos. With




Figure 10a: Scatter-plot showing the AMANDA-reconstructed θ -angle of atmospheric muons versus the MC, at several cut levels.




Figure 10b: $\theta_{\text{rec}} - \theta_{\text{MC}}$ for reconstructed atmospheric Monte Carlo simulated muons with at least five residuals in the interval $[-15; 15]$ ns.




Figure 11: Atmospheric neutrino events passing the cuts imposed on the data to search for neutrinos from WIMP annihilation in the center of the earth.

standard statistical techniques this result can be converted into an upper limit on an excess flux of WIMP origin; see Fig. 12.

In order to interpret this result, we have simulated AMANDA-80 sensitivity to the 2 dominant WIMP annihilation channels[16]: into $b\bar{b}$ and W^+W^- . The upper limits on the WIMP flux are shown in Fig. 12 as a function of the WIMP mass. Limits below 100 GeV WIMP mass are poor because the neutrino-induced muons (with typical energy $\simeq m_\chi/6$) fall below the AMANDA-80 threshold. For the heavier masses, limits approach the limits set by other experiments in the vicinity of $10^{-14} \text{ cm}^{-2} \text{ s}^{-1}$. We have previously discussed how data, already on tape from AMANDA-300, will make new incursions into the parameter space of supersymmetric models shown in Fig. 1.

Event ID#	4706879	8427905
α [m/ns]	0.19	0.37
Length [m]	295	182
Closest approach [m]	2.53	1.23
$\theta_{rec} [^\circ]$	14.1	4.6
$\phi_{rec} [^\circ]$	92.0	348.7
Likelihood/OM	5.9	4.2
OM multiplicity	14	8
String multiplicity	4	2

Table 1: Characteristics of the two events reconstructed as up-going muons.




Figure 12: Upper limit at the 90% confidence level on the muon flux from the center of the earth as a function of neutralino mass. The light shaded band represents the W^+W^- annihilation channel and the dark one represents the $b\bar{b}$ annihilation channel. The width of the bands reflects the inadequate preliminary calculation.

Acknowledgements

The AMANDA collaboration is indebted to the Polar Ice Coring Office and to Bruce Koci for the successful drilling operations, and to the National Science Foundation (USA), the Swedish National Research Council, the K. A. Wallenberg Foundation and the Swedish

Polar Research Secretariat. F.H. was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

References

- [1] For a review, see T.K. Gaisser, F. Halzen and T. Stanev, *Phys. Rep.* **258**(3), 173 (1995); R. Gandhi, C. Quigg, M. H. Reno and I. Sarcevic, *Astropart. Phys.*, **5**, 81 (1996).
- [2] The Pierre Auger Project Design Report, Fermilab report (Feb. 1997) and references therein.
- [3] F. Halzen, *The case for a kilometer-scale neutrino detector*, in Nuclear and Particle Astrophysics and Cosmology, Proceedings of Snowmass 94, R. Kolb and R. Peccei, eds.
- [4] S. Scopel, *Proceedings of the International Workshop on Aspects of Dark Matter in Astrophysics and Particle Physics*, Heidelberg, Germany, (1996); see also L. Bergström, J. Edsjo and P. Gondolo, *Phys. Rev. D* (to be published).
- [5] O. Suvorova et al., *Baksan neutralino search*, presented at the First International Workshop on Non-Accelerator New Physics (NANP97), Dubna, Russia, July 7–12, (1997).
- [6] G. Jungman, M. Kamionkowski and K. Griest, *Phys. Rep.* **267**, 195 (1996).
- [7] P. Belli, these proceedings.
- [8] S. W. Barwick *et al.*, *The status of the AMANDA high-energy neutrino detector*, in Proceedings of the 25th International Cosmic Ray Conference, Durban, South Africa (1997).
- [9] S. Tilav *et al.*, *First look at AMANDA-B data*, in Proceedings of the 25th International Cosmic Ray Conference, Durban, South Africa (1997).
- [10] S. W. Barwick *et al.*, *Proceedings of the 22nd International Cosmic Ray Conference*, Dublin (Dublin Institute for Advanced Studies, 1991), Vol. 4, p. 658.
- [11] The AMANDA collaboration, *Science* **267**, 1147 (1995).
- [12] C. Wiebusch *et al.*, *Muon reconstruction with AMANDA-B*, in Proceedings of the 25th International Cosmic Ray Conference, Durban, South Africa (1997).
- [13] T. Miller *et al.*, *Analysis of SPASE-AMANDA coincidence events*, in Proceedings of the 25th International Cosmic Ray Conference, Durban, South Africa (1997).
- [14] A. Bouchta, University of Stockholm, PhD thesis (1998)
- [15] J. E. Jacobsen, University of Wisconsin, PhD thesis (1996)
- [16] Muflux code written by J. Edsjoe, implemented as an event generator by A. Goobar; see also L. Bergström, J. Edsjö and P. Gondolo, *Indirect neutralino detection rates in neutrino telescopes*, *Phys. Rev. D* **55** 1765, (1997).